



UNIVERSITI PUTRA MALAYSIA

**EVALUATION OF SOLID STATE FERMENTATION BY ASPERGILLUS
NIGER TO IMPROVE THE NUTRITIVE VALUE OF PALM KERNEL
CAKE FOR BROILERS**

KHIN HNIN SWE.

FP 2004 2

**EVALUATION OF SOLID STATE FERMENTATION BY *ASPERGILLUS
NIGER* TO IMPROVE THE NUTRITIVE VALUE OF PALM KERNEL
CAKE FOR BROILERS**

KHIN HNIN SWE

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2004



**EVALUATION OF SOLID STATE FERMENTATION BY *ASPERGILLUS NIGER*
TO IMPROVE THE NUTRITIVE VALUE OF PALM KERNEL CAKE FOR
BROILERS**

**Disertai CD-ROM / disket yang boleh diperolehi
di Bahagian Media dan Arkib
(*Accompanying CD-ROM / disk available
at the Media and Archives Division*)**

By

KHIN HNIN SWE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

December 2004



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

EVALUATION OF SOLID STATE FERMENTATION BY *ASPERGILLUS NIGER* TO IMPROVE THE NUTRITIVE VALUE OF PALM KERNEL CAKE FOR BROILERS

By

KHIN HNIN SWE

December 2004

Chairman : Professor Abd. Razak Bin Alimon, Ph.D.

Faculty : Agriculture

A study was conducted to examine the effects of solid-state fermentation (SSF) by *Aspergillus niger* on the nutrient contents of palm kernel cake (PKC). The effects of different types and levels of alcohols were investigated on the enzyme activity and nutrient contents of fermented PKC+ rice bran (RB). The results showed that the neutral detergent fibre (NDF) and acid detergent fibre (ADF) contents of fermented substrate (FS) (PKC+RB) were decreased by 47.3% and 22.6% respectively, while the crude protein (CP) was increased by 28.7% after 8 days of fermentation and the activity of mannanase was 109.7 U/g at 6 days of fermentation. There were no significant effects ($P>0.05$) of ethanol and methanol on the nutrient contents and citric acid production in the fermented substrate.

An experiment was conducted to improve the fermentation process of *A. niger* on PKC. Rice bran was added to PKC in varying proportions. Three different proportions of rice bran (10%, 20% and 30% of total substrate) were used and the fermentation was carried out for 8 days. The results showed that the nutrient contents of FS were not significantly ($P>0.05$) affected by the RB added. Nevertheless, among the different substrate groups used, 70% PKC+ 30% RB showed the lowest concentration of ochratoxin (43.3ppb). The ochratoxin level of 100% PKC substrate group was 49.8 ppb. Therefore, based on the ochratoxin level, the substrate group consisting of 70% PKC + 30% RB was used as the substrate for SSF. The fermented substrate was used as an ingredient in the broiler ration.

Four experimental diets were used in the feeding trial. They were: 1) control diet (without PKC), 2) diet containing 20% PKC, 3) diet containing 20% FS and 4) diet containing 25% FS. A total of 60 birds, 15 birds per treatment, 3 weeks old Arbor Acre were used. The birds were placed in individual cages. The growth trial was continued up to 42 days. The results showed poor ($P<0.05$) growth performances of birds fed diets containing FS and the best performance was recorded in birds fed control diet. There were no significant ($P>0.05$) differences in the performance of birds fed diet containing 20% PKC and those fed control diet. Cumulative FCR of birds fed control diet was 1.85 whereas that of birds fed dietary treatment 4 was 2.42.

The study on histopathological examination of visceral organs of chickens fed diet containing FS showed some lesions of ochratoxicosis where the immune organ (bursa

of Fabricius) was adversely affected. The lesions in liver were less than those of the kidney.

An attempt was carried out to delay sporulation of fungal culture. In this study, 14% ammonium sulphate was observed as the best concentration to add in culture during SSF of substrate (PKC+RB) by *A. niger* to delay sporulation. The digestibility of fibre was improved by adding ammonium sulphate in the culture.

Although the nutritive value of fermented PKC was enhanced by SSF with *A. niger*, the feeding trial carried out in broilers showed no beneficial effect of using the fermented substrate as a ration component in poultry feed. Nevertheless, the information obtained in this study would be beneficial in the understanding of the biochemical changes that occur in PKC during SSF with *A. niger*.

Abstrak yang dikemukakan kepada Senat Universiti Putra Malaysia untuk memenuhi keperluan bagi ijazah Doktor Falsafah.

**PENILAIAN FERMENTASI DALAM BENTUK PEPEJAL HAMPAS KELAPA
OLEH *ASPERGILLUS NIGER* UNTUK MENINGKATKAN NILAI NUTRITIF
HAMPAS KELAPA SAWIT UNTUK AYAM PEDAGING**

Oleh

KHIN HNIN SWE

Disember 2004

Pengerusi : Profesor Abd. Razak Bin Alimon, Ph.D.

Fakulti : Pertanian

Satu kajian telah dijalankan untuk menilai kesan fermentasi dalam bentuk pepejal oleh *Aspergillus niger* ke atas kandungan nutrien dari hampas kelapa sawit.. Dalam kajian pertama, kesan dari jenis dan tahap kandungan alkohol yang berbeza telah dikaji ke atas aktiviti enzim dan kandungan nutrien dari hampas padi + PKC yang telah difermentasikan. Keputusan telah menunjukkan kandungan NDF dan ADF dari substrat yang difermentasi (FS) (PKC+RB) adalah menurun sebanyak 47.3% dan 22.6%, sementara protein kasar meningkat 28.7% selepas 8 hari fermentasi dan aktiviti mannanase adalah 109.7 U/g pada fermentasi selama 6 hari. Tiada perbezaan signifikan ($P>0.05$) kesan dari ethanol dan methanol pada kandungan nutrien dan penghasilan asid sitrik dalam substrat telah difermentasi.

Kajian telah dijalankan untuk menambahbaik proses fermentasi oleh *A. niger* ke atas PKC. Hampas padi telah ditambah kepada PKC dalam kadar yang berbeza. Tiga kadar hampas padi (10%, 20% dan 30% dari jumlah substrat) telah digunakan dan fermentasi telah dilakukan selama 8 hari. Keputusan telah menunjukkan bahawa kandungan nutrien dari FS adalah tambahan hampas padi tiada perbezaan yang signifikan. Walaupun begitu, antara kumpulan substrat yang digunakan, 70% PKC + 30% RB menunjukkan kepekatan oktratoksin (43.3ppb) yang terendah. Oktratoksin pada kumpulan tahap substrat PKC 100% adalah 49.8 ppb. Oleh itu, berdasarkan kepada tahap oktratoksin kumpulan substrat yang mengandungi 70% PKC + 30% RB telah digunakan sebagai substrat untuk SSF. Substrat yang difermentasi telah digunakan sebagai bahan kandungan dalam ransum ayam daging.

Empat ujian diet telah digunakan dalam pengujian pemakanan. Mereka adalah: 1) diet kawalan (tanpa PKC), 2) diet mengandungi 20% PKC, 3) diet mengandungi 20% FS dan 4) diet mengandungi 25% FS. Sejumlah dari 60 ekor ayam Arbor Acre, 15 ekor setiap kumpulan, dan berumur 3 minggu digunakan. Ujian tumbesaran dilakukan dalam sangkar berasingan sehingga berumur 42 hari. Keputusan menunjukkan tumbesaran adalah rendah untuk ayam yang diberi diet FS dan terbaik adalah kumpulan ayam yang diberi diet kawalan. Tiada perbezaan yang signifikan ($P>0.05$) dalam tumbesaran ayam yang diberi makan diet mengandungi 20% PKC dan yang diberi makanan diet kawalan. FCR terkumpul dari ayam yang diberi makan diet kawalan. FCR terkumpul dari ayam yang diberi makan diet kawalan adalah 1.85 sementara ayam yang diberi diet 4 adalah 2.42.

Kajian ke atas histopatologi terhadap organ visceral ayam yang diberi diet mengandungi FS menunjukkan kerosakan oktratoksikosis di mana bursa Fabricius menunjukkan kesan sebaliknya. Kerosakan pada hati adalah kurang berbanding dengan buah pinggang

Kajian telah dijalankan untuk menanggukkan sporulisasi kultur kulat. Dalam kajian ini, 14% ammonium sulfat dicatatkan sebagai kepekatan terbaik untuk ditambahkan dalam kultur sewaktu SSF dari substrat (PKC +RB) oleh *A. niger* untuk penanguhan sporulisasi. Pencernaan tenaga dan gentian adalah meningkat dengan penambahan ammonium sulfat pada kultur.

Walaupun nilai nutritif dan PKC difermentasi adalah ditingkatkan oleh SSF dengan *A. niger*, percubaan pemakanan dalam ayam daging menunjukkan tiada kesan baik dari penggunaan bahan difermentasi sebagai bahan ration makanan ayam. Walaupun begitu, maklumat yang telah diperolehi dari kajian ini akan memanfaatkan dalam kefahaman perubahan yang berlaku dalam PKC semasa SSF dengan *A. niger*.

ACKNOWLEDGEMENTS

I would like to express my sincere appreciation and deepest gratitude to Professor Dr. Abdul Razak Alimon, Chairman of the supervisory committee for his invaluable guidance, advice, continuous supervision and encouragement throughout this study and in the preparation of this thesis.

I wish to express my heartfelt thankfulness to Associate Professor Dr. Norhani bt. Abdullah and Dr. Mohd Jaafar Daud, Senior Research Officer of MARDI members of the supervisory committee for their valuable advice, patience, help and encouragement during this study and in the preparation of this thesis.

Deep appreciation is also expressed to my sponsor, MTCP (Malaysian Technical Cooperation Programme) for providing me a scholarship to pursue this doctoral study.

I am also much indebted to Minister of Agriculture, Malaysia and Minister of Livestock and Fisheries of Myanmar for offering the opportunity to study in Malaysia. I am also grateful to the Director General of respective departments and Rector/ dean of both countries for their important roles in materializing the scholarship programme for me.



I would like to acknowledge to Ms. Noraini Samat, Research Officer of MARDI. Sincere appreciation and gratefulness are also extended to Ms. Sarah Rasol, Assistant Research Officer and all staff of Bioprocess laboratory from MARDI for their technical assistance and kindness. A special thanks is due to Mr. Mohd Sharudin Bin Mohd Ali for helping me tireless transportation to MARDI. My thanks are in particular to Mr. Ibrahim Mohsin, Mr. Saparin Denim, Mr. Hj. Abdul Halin B. Hj. Mohd. Isa and Mr. Bakari Abd. Rahman for helping me during the laboratory work in Universiti Putra Malaysia. My appreciation is especially to Mr. Ismail, Department of Veterinary Service in Petaling Jaya for his enthusiastic cooperation during amino acid assay. I would like to thank Zawawi Ismail and Lai Pui Wah, postgraduate students of Animal Science department for helping me to write the abstract of thesis in Bahasa Melayu.

Deepest thanks is due to Rector Dr. Aung Than, University of Veterinary Science Myanmar for allowing me study leave and his shouldering on all my duties while I was abroad.

My appreciations are also due to my father, my mother, my sisters and brothers for their moral support and love.

Last but not least, very special thanks go to my husband, Associate Professor Dr. Myint Wai, University of Veterinary Science Myanmar and lovely daughter Honey Hnin Wai for their moral support, encouragement, patience, love and understanding over the many selfish times for completing this study.

I certify that an Examination Committee met on 16th December 2003 to conduct the final examination of Khin Hnin Swe on her Doctor of Philosophy thesis entitled "Evaluation of Solid State Fermentation By *Aspergillus niger* to Improve the Nutritive Value of Palm Kernel Cake for Broilers" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Zulkifli Idrus, Ph.D.

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Liang Juan Boo, Ph.D.


Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Azhar Kasim, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Hyung Tai Shin, Ph.D.

Professor
Faculty of Life Science & Technology
Sungkyunkwan University
Korea
(Independent Examiner)



ZAKARIAH ABD. RASHID, Ph.D.
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 JAN 2005

This thesis is submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Abdul Razak Alimon, Ph.D.

Professor

Faculty of Agriculture

University Putra Malaysia

(Chairman)

Norhani bt. Abdullah, Ph.D.

Associate Professor

Faculty of Science and Environmental studies

Universiti Putra Malaysia

(Member)

Mohd. Jaafar Daud, Ph.D.

Senior Research Officer

Livestock Research Centre

Malaysian Agricultural Research and Development Institute

(Member)



AINI IDERIS, Ph.D.

Professor / Dean


School of Graduate Studies

Universiti Putra Malaysia

Date: 28 JAN 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



KHIN HNIN SWE

Date: 5 / 2 / 05 .

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVALS	x
DECLARATION	xii
LIST OF TABLES	xvii
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxii
 CHAPTER	
 1. GENERAL INTRODUCTION	 1
 2. LITERATURE REVIEW	 6
2.1 Introduction	6
2.2 Palm Kernel Cake	6
2.2.1 Energy in PKC	9
2.2.2 Protein in PKC	10
2.2.3 Other nutrients in PKC	12
2.3 Definition of solid state fermentation	12
2.3.1 History and Development of SSF	13
2.3.2 General consideration about solid-state fermentation	16
2.3.2.1 Temperature	16
2.3.2.2 pH	17
2.3.2.3 Moisture and water activity	18
2.3.2.4 Aeration	20
2.3.2.5 Substrate concentration	21
2.3.3 Scaling-up	21
2.4 Growth kinetics of filamentous fungi	23
2.4.1 Biomass in solid state fermentation	26
2.4.2 Organic acid produced by SSF	26
2.4.3 Factors affecting sporulation	31
2.4.3.1 Effect of nutrition	32
2.4.3.2 Other factors affecting sporulation	33
2.4.4 Mycotoxin produced by filamentous fungi	36
2.5 Nutrient requirements of poultry	39
2.5.1 Palm Kernel Cake in broiler and layer ration	42
 3. GENERAL MATERIALS AND METHODS	 45
3.1 Fermentation technique	45
3.1.1 Inoculum preparation (Spore suspension)	45
3.1.2 Substrate preparation	45



3.1.3 Powder inoculum preparation	46
3.1.4 Mineral solution	46
3.1.5 Solid state fermentation	46
3.2 Preparation of crude filtrate of fermented substrate	47
3.2.1 Dinitrosalicylic acid method for measuring total reducing sugar	47
3.2.2 Mannanase activity assay	48
3.2.3 Bio-rad assay method for soluble protein determination	51
3.3 Proximate Analysis	52
3.3.1 Determination of moisture and dry matter	52
3.3.2 Determination of ash	53
3.3.3 Determination of neutral detergent fibre	53
3.3.4 Determination of acid detergent fibre	54
3.3.5 Determination of ether extract	55
3.3.6 Determination of crude protein	56
3.4 Determination of minerals	57
3.5 Determination of amino acids	58
3.6 Procedure for TME (True Metabolizable Energy) determination	62
3.7 Biomass determination	64
3.8 Citric acid determination	66
3.9 Determination of organic carbon	68
3.10 The procedure of Scale-up fermentation	70
3.11 Enzyme linked immunoassay for the quantity analysis of ochratoxin	71
4. EVALUATION OF SSF BY <i>ASPERGILLUS NIGER</i> ON THE NUTRITIVE VALUE OF PKC+RB WITH OR WITHOUT ADDITION OF ALCOHOL	73
4.1 Introduction	73
4.2 Materials and Methods	75
4.2.1 Chemical analysis	76
4.2.2 Statistical Analysis	76
4.3 Results	76
4.3.1 Effects of alcohol	76
4.3.2 Fermentation time	79
4.3.3 The effect of alcohol at different fermentation times	82
4.4 Discussion	84
4.5 Conclusion	90
5. THE EFFECTS OF VARYING LEVELS OF RICE BRAN ON THE FERMENTATION OF PKC	92
5.1 Introduction	92
5.2 Materials and Methods	93
5.2.1 Chemical analysis	94
5.2.2 Statistical analysis	95
5.3 Results	95
5.4 Discussion	107

5.5 Conclusion	114
6. THE EFFECT OF FERMENTED PKC ON BROILER PERFORMANCE	116
6.1 Introduction	116
6.2 Materials and Methods	118
6.2.1 Animals and housing and management	118
6.2.2 Diets	119
6.2.3 Fermentation procedure	119
6.2.4 Digestibility trial	120
6.2.5 Chemical analysis	120
6.2.6 Statistical Analysis	122
6.3 Results	122
6.3.1 Live-weights	122
6.3.2 Feed intake	123
6.3.3 Weight gain	123
6.3.4 Feed Conversion Ratio (FCR)	125
6.3.5 Relative organ weight and Carcass characteristics	125
6.3.6 Digestibility	126
6.4 Discussion	130
6.5 Conclusion	134
7. HISTOPATHOLOGICAL EXAMINATION OF VISCERAL ORGANS OF CHICKEN FED DIET CONTAINING FERMENTED PKC	135
7.1 Introduction	135
7.2 Materials and Methods	137
7.2.1 Experimental birds and diets	137
7.2.2 Procedure	137
7.2.3 Scoring of lesions	138
7.3 Results	140
7.4 Discussion	150
7.5 Conclusion	152
8. THE EFFECT OF AMMONIUM SULPHATE ON THE NUTRITIVE VALUE OF FERMENTED PKC BY <i>ASPERGILLUS NIGER</i>	153
8.1 Introduction	153
8.2 Materials and Methods	155
8.2.1 Experiment 1	155
8.2.1.1 Chemical Analysis	156
8.2.1.2 CFU test	156
8.2.1.3 Statistical Analysis	157
8.2.2 Experiment 2	157
8.2.2.1 Chemical Analyses	158
8.2.2.2 Statistical Analysis	158
8.3 Results	159
8.3.1 Experiment 1	159

8.3.1.1 Effect of levels of ammonium sulphate	159
8.3.1.2 Effect of different incubation periods	159
8.3.1.3 The development of spore in the culture	162
8.3.1.4 The effect of ammonium sulphate on CFU, NDF, ADF values of FS at different incubation periods	166
8.3.2 Experiment 2	171
8.3.2.1 Effect of ammonium sulphate on nutrient contents and Mannanase activity	171
8.3.2.2 Effect of incubation periods	171
8.3.2.3 Effect of ammonium sulphate on the parameters measured at two incubation periods	173
8.4 Discussion	183
8.5 Conclusion	189
9 GENERAL DISCUSSION	190
10 GENERAL CONCLUSIONS	199
REFERENCES	201
APPENDICES	222
BIODATA OF THE AUTHOR	226

LIST OF TABLES

Table	Page
1 Chemical composition of palm kernel cake (% DM basis)	8
2 History and development of solid state fermentation	15
3 Important mycotoxins and the producer fungi	37
4 Growth and fed consumption of broiler chickens	40
5 The enzyme activities of each dilution for standard curve	50
6 The mean value of NDF, ADF, CP and soluble protein content, mannanase activity, reducing sugar of fermented substrate, glucosamine content of fungus and citric acid content of the culture in different treatments	78
7 The mean values of NDF, ADF, CP and soluble protein, mannanase activity, reducing sugar of fermented substrate, glucosamine content of fungus and citric acid of the culture at various fermentation periods	81
8 The mean value of mannanase activity of culture for each treatment at different fermentation times (U/g)	83
9 The mean value of glucosamine contents of fungus for each treatment at different fermentation time ($\mu\text{g/g}$)	84
10 The percentage of DM, CP and ash of different substrates at day 0 and day 8 fermentation	96
11 The percentage of ether extract, NDF and ADF of substrate at day 0 and day 8 fermentation	98
12 Reducing sugar content of untreated and fermented substrate at day 0 and day 8 fermentation	99
13 Mannanase activity, glucosamine content of fungus and citric acid of culture for different substrate at 8 days fermentation	100
14 Gross energy and TME value of unfermented and fermented substrate for two treatments at day 0 and 8 fermentation	101



15	Carbon and nitrogen ratio of different substrate groups	102
16	Amino acids profile of substrate at 0 day (% DM basis)	104
17	Amino acids profiles of fermented substrates at 8 days fermentation (%DM basis)	105
18	Amino acids profiles of substrates for all treatments at day 0 and 8 days fermentation (%DM basis)	106
19	The effect of different substrate groups on ochratoxin level of fermented substrate for different treatments	107
20	Composition of the diets using untreated PKC and fermented substrate	121
21	Growth performance of broilers fed different diets for three weeks (Mean \pm SE)	124
22	Organs and abdominal fat pad weights expressed as percentages of live-weight (Mean \pm SE)	126
23	Carcass characteristics of broilers fed experimental diets	127
24	Digestibility of dry matter, CP, NDF, ADF, Ca and phosphorous of experimental diets	128
25	The concentration of ochratoxin for each treatment and scoring of lesions for bursa of Fabricius, kidney and liver of chickens	140
26	The effects of levels of ammonium sulphate and incubation periods on CFU value of culture, NDF & ADF contents of FS	161
27	The effect of ammonium sulphate on CFU (colony forming unit) value of culture	166
28	The effect of different levels of ammonium sulphate on the NDF of fermented substrate at different incubation periods	168
29	The effect of different levels of ammonium sulphate on ADF contents of fermented substrate at different incubation periods	169
30	The effects of ammonium sulphate and incubation times on NDF, ADF, CP, ether extract, ash content, reducing sugar of FS and mannanase	172

activity of culture

31	The effect of ammonium sulphate on NDF, ADF, CP, Ether extract and ash of FS at two fermentation times	174
32	The effect of ammonium sulphate on Reducing sugar level of FS at different fermentation times	178
33	The effect of ammonium sulphate on Mananase enzyme activity of culture at different fermentation times	178
34	The effect of ammonium sulphate on Gross energy of FS at different fermentation times	180
35	The effect of ammonium sulphate on TME of FS at different fermentation times	181
36	The effect of ammonium sulphate on apparent digestibility of NDF of FS	182
37	The effect of ammonium sulphate on ochratoxin level (ppb) of fermented substrate	182

LIST OF FIGURES

Figure		Page
1	Phases of fungal growth	24
2	Metabolic process associated with the production of citric acid and its conversion to other other products	30
3	Standard curve of glucose	48
4	Standard curve of mannanase activity	50
5	Standard curve of soluble protein	51
6	Standard curve of glucosamine	66
7	Standard curve of citric acid	68
8	Reducing sugar and soluble protein of FS	88
9	The glucosamine content of fungus and mannanase activity	89
10	Glucosamine content of fungus and citric acid production	91
11	Digestibility of nutrients for each experimental diet	129
12	Scoring of lesions for bursa of Fabricius in each treatment	141
13	Micrograph of liver (T1)	143
14	Micrograph of liver (T2)	143
15	Micrograph of liver (T3)	143
16	Micrograph of liver (T4)	143
17	Micrograph of kidney (T1)	146
18	Micrograph of kidney (T2)	146
19	Micrograph of kidney (T3)	146

20	Micrograph of kidney (T4)	146
21	Micrograph of bursa of Fabricius (T1)	148
22	Micrograph of bursa of Fabricius (T2)	149
23	Micrograph of bursa of Fabricius (T3)	149
24	Micrograph of bursa of Fabricius (T4)	149
25	Photograph of FS with or without ammonium sulphate at 3 days fermentation	163
26	Photograph of FS with or without ammonium sulphate at 4 days fermentation	164
27	Photograph of FS with or without ammonium sulphate at 5 day fermentation	165
28	The effect of ammonium sulphate on CFU of culture	167
29	The effect of different levels of ammonium sulphate on the NDF content of fermented substrate	170
30	The effect of different levels of ammonium sulphate on the ADF contents of the fermented substrate	170
31	The effect of 14% ammonium sulphate on the NDF content of FS	175
32	The effect of 14% ammonium sulphate on the ADF content of FS	175
33	The effect of 14% ammonium sulphate on the CP content of FS	176
34	The effect of 14% ammonium sulphate on the EE contents of FS	176

LIST OF ABBREVIATIONS

AABA	- L- α -Amino-n-butyric acid
ADF	- Acid Detergent Fibre
AME	Apparent metabolizable energy
Ca	- Calcium
CFU	- Colony Forming Unit
C: N	- Carbon Nitrogen Ratio
CP	- Crude Protein
C/P ratio	- Calorie Protein ratio
DM	- Dry Matter
EE	- Ether Extract
FCR	- Food Conversion Ratio
FS	- Fermented substrate (PKC+RB)
GE	- Gross Energy
HPLC	- High Performance Liquid Chromatography
ISR	Internal Standard Ratio
LD ₅₀	- Lethal Dose at 50%
MBTH	- 3-methyl-2-benzothiozalone hydrochloride
NDF	- Neutral Detergent Fibre
NRC	- National Research Council
NS	Not Significantly Different
NSPs	- Non Starch polysaccharides

OA	Ochratoxin A
P	- Phosphorous
PDA	- Potato Dextrose Agar
PKC	- Palm Kernel Cake
PKM	- Palm Kernel Meal
POME	- Palm Oil Mill Effluent
POS	Palm Oil Sludge
PPF	- Palm Press Fibre
RB	Rice Bran
SD	Standard Deviation of t-test
SDW	Sample Dry Weight
SPSS	- Statistical Package for Social Scientist
SSF	- Solid-State Fermentation
TME	- True Metabolizable Energy